

LEVELINA SOLUTION

COMPOSITION

Per 100ml

Bifonazol (D.C.I) 1g

Excipients: Ethanol, isopropyl myristate

PHARMACEUTICAL FORM AND CONTENTS OF PACKET

Levelina solution, 30ml bottle

ACTION

LEVELINA (Bifonazol) is a broad spectrum antimycotic which deals with dermatophytes, yeasts, mold and other fungal infections such *Malessezia furfus*; it is also effective against *Corynebacterium minutissimum*. Action against dermatophytes (e.g. *Trichophyton* spp.) is fundamentally fungicidal in vitro, and in yeasts it is strongly fungistatic.

Name of person or entity authorized for commercialization and production of this product:

Laboratorios ERN, S.A. Pedro IV, 499-08020 Barcelona, España

INDICATIONS

Skin mycosis caused by dermatophytes, yeasts, molds, and other fungi, such as *Malessezia furfur* and *Corynebacterium minutissimum*. These could be, for example; Mycosis of the feet and hands (e.g. athlete's foot, ringworm of the hands) Mycosis of the body and cutaneous complaints (e.g. ringworm), *Pitriasis versicolor* and *Erythrasma*.

COUNTERINDICATIONS

Hypersensitivity to Bifonazol.

INCOMPATIBILITIES

None have been described.

WARNING

Frequent applications may cause dry skin and irritation.

POSODOGY

Unless advised otherwise by a doctor: Levelina (Bifonazol) is applied once a day, preferably before going to bed, spreading a small amount of solution on the affected area. Generally it is sufficient to apply a few drops for an area the size of your hand. To hygienically support the treatment it is advisable to carefully wash and dry the affected area. To obtain lasting results treatment with LEVELINA (Bifonazol) should not be interrupted after the symptoms have disappeared, but should be continued for a determined period of time:

Interdigital mycosis of the foot and athlete's foot: 3 weeks

Mycosis of the body, hands and cutaneous complaints (ringworm etc.): 2-3 weeks

Pitiriasis versicolor and Erythrasma: 2 weeks

Superficial cutaneous candidiasis: 2-4 weeks

OVERDOSE AND ITS TREATMENT

Given the concentration of the active principle and the form of administration, intoxication is not possible even in the case of accidental ingestion.

In the event of overdose or accidental ingestion consult El Servicio de Información Toxicológica. Tel 91 562 04 20.

ADVERSE REACTIONS

In rare cases light cutaneous reactions may appear. Any adverse reaction which is not described in the prospectus should be reported to your doctor.

EXPIRY

This medicine should not be used after the expiry date indicated on the packaging.

PRESCRIPTION ONLY

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN.

TITULAR AUTHORIZED FOR COMMERCIALIZATION

Laboratorios ERN, S.A. Pedro IV, 499-08020 Barcelona, España

RESPONSIBLE FOR MANUFACTURING

Laboratorios ERN, S.A. Polígono Industrial Can Salvatella, c/ Gorcs i Lladó, 188 Barberá de Vallés. Barcelona. España

Laboratorios ERN, S.A. c/ Perú, 228 - 08020 Barcelona. España