

# Aftajuventus

## Local treatment of aphthas and ulcerations in the buccal mucous membrane

### COMPOSITION

Hydrocortisone hemisuccinate (ICD).....60 mg  
Benzalkonium chloride (ICD).....3 mg  
Excipient: glycerine, q.s. ....30 mg

### INDICATIONS

Aphthas  
Infectious stomatitis (mouth rot)  
Catarrhal stomatitis  
Muguet

### DOSAGE

2 or 3 applications a day on the affected area, making use of the stick-spatula (always according to the practitioner's prescription).

### GUIDELINES FOR ITS CORRECT USE

After every use and before putting the stick-spatula into the bottle to get more liquid, it is advisable to clean the stick with cotton soaked in alcohol.

### CONTRAINDICATIONS AND PRECAUTIONS

Treatment should not be prolonged for long periods nor for herpes-type diseases.

### INCOMPATIBILITIES

It has no incompatibilities.

### INTERACTIONS

Interactions have not been described in the topical use of this formula.

### SIDE EFFECTS

Its prolonged application for weeks may cause symptoms of hypercorticism, which disappear when use is suspended.

### INTOXICATION AND ITS TREATMENT

The ingestion of one container does not cause toxic effects, although the age and state of the patient must be considered in each case.

In case of overdose or accidental ingestion, immediately consult the Toxicology Information Service, telephone 915620420.

### FORMAT

Bottle with applicator, containing 30 ml.

### WITH MEDICAL PRESCRIPTION

### MEDICINES MUST BE KEPT OUT OF REACH OF CHILDREN

Holder of the authorisation for the commercialisation and fabrication of the product.  
LABORATORIOS ERN, S.A. Pedro IV, 499 - 08020 Barcelona, España

Tel: 93 314 80 11 Fax: 93 314 40 96

Responsible for fabrication.

Laboratorios ERN, S.A. Gorcs i LLadó, 188 – 08210 Barberá del Vallés, Barcelona.  
España.

Laboratorios ERN, S.A. Perú, 228 – 08020 Barcelona. España.